

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
20 December 2001 (20.12.2001)

PCT

(10) International Publication Number
WO 01/95907 A1(51) International Patent Classification⁷: A61K 31/428,
A61P 25/00

(21) International Application Number: PCT/IL01/00534

(22) International Filing Date: 12 June 2001 (12.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
136687 12 June 2000 (12.06.2000) IL(71) Applicant (for all designated States except US): MOR -
RESEARCH APPLICATIONS LTD. [IL/IL]; Hasivim
Street 23, Kiryat Matalon, 49517 Petach Tikva (IL).


(72) Inventors; and

(75) Inventors/Applicants (for US only): MELAMED, Eldad
[IL/IL]; Tagor Street 44, Ramat Aviv, 69341 Tel Aviv (IL).
OPHEN, Daniel [IL/IL]; 38955 Kfar Haroeh (IL).(74) Agent: REINHOLD COHN AND PARTNERS; P.O.
Box 4060, 61040 Tel Aviv (IL).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,
ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**Published:**

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PHARMACEUTICAL COMPOSITION FOR THE TREATMENT OF MULTIPLE SCLEROSIS



(57) Abstract: Riluzole, a drug that inhibits glutamathergic release, is shown to be effective in the prevention and treatment of multiple sclerosis (MS). The effect of Riluzole is shown in an animal model of MS, an experimental autoimmune encephalomyelitis (EAE) model produced by injection of myelin oligodendrocyte glycoprotein (MOG) to animals. Administration of Riluzole to such animals before they develop the MS-related symptoms markedly reduced the incidence and clinical severity of the disease in such animals. Moreover, treatment of such animals after the appearance of severe MS-related symptoms, also markedly slowed down the progression of the disease and improved the clinical manifestations.

Applicants: Daniella Licht et al.
Serial No.: 10/772,911
Filed: February 5, 2004
Exhibit 4

BEST AVAILABLE COPY

WO 01/95907 A1

PHARMACEUTICAL COMPOSITION FOR THE TREATMENT OF MULTIPLE SCLEROSIS

FIELD OF THE INVENTION

This invention concerns pharmaceutical compositions for the treatment of
5 multiple sclerosis (MS).

BACKGROUND OF THE INVENTION

Multiple Sclerosis (MS) is a disorder of the central nervous system, involving decreased nerve function associated with the formation of scars on the myelin covering of nerve cells. This disease affects approximately 1 out of 1,600
10 people, with women being affected 60% of the time. The disorder most commonly begins between the ages of 20 to 40, and is one of the major causes of disability in adults under age 65.

Multiple sclerosis involves repeated episodes of inflammation of nervous tissue in any area of the central nervous system, including the brain and spinal cord.
15 The location of the inflammation varies from one person to another and from episode to episode. The inflammation destroys the myelin sheath covering the nerve cells in that area, which causes the formation of multiple areas of scar tissue (sclerosis) along the covering of the nerve cells. Sclerosis slows or blocks the transmission of nerve impulses in that area, resulting in the development of the
20 symptoms of MS.

Symptoms vary considerably, since the location and extent of each attack varies. There is usually a stepwise progression of the disorder. At the initial stages ("relapsing-remitting" stage) the episodes of onset of symptoms last days, weeks or months, alternating with times of reduced or no symptoms (remission) and periods
25 of recurrence (relapse). During relapse there is an appearance of a new symptom, the reappearance of a previous symptom or the worsening of an existing symptom. At more advance stages (termed: "chronic-progressive" stage which may be either

- 2 -

primary or secondary), there is a progressive deterioration of nerve function which is probably caused by the irreversible destruction of nerve axons.

The exact cause of the inflammation associated with MS is unknown. Several geographic studies indicate that there may be an environmental factor involved with MS. There seems to be a familial tendency toward the disorder, with a higher incidence in certain family groups than in the general population indicating a possible genetic involvement. An increase in the number of immune cells in the body of a person with MS indicates that there may be a type of immune response that triggers the disorder.

10 The most frequent theories about the cause of multiple sclerosis include: infection by a virus-type organism; an abnormality of the genes responsible for control of the immune system; or a combination of both factors.

There is no known cure for multiple sclerosis and treatment is aimed at controlling systems and maintaining functions to give the patient a maximum quality of life.

15 Medications vary depending on the symptoms that occur. Baclofen, dantrolene, diazepam or other anti-spasmodic medications are used to reduce muscle spasticity. Cholinergic medications may be helpful to reduce urinary problems. Antidepressant medications may be helpful for mood or behavior symptoms. Amantadine may be given for fatigue.

Corticosteroids or ACTH are frequently used to suppress the inflammation in an attempt to reduce the duration of an attack. Medications that suppress the immune are also often used. Recently it has been found that Interferon may also be helpful for some people.

25 Recent studies (Pitt *et al.*, *Nature Medicine*, 6(1), 67-70, 2000; Smith *et al.*, *Nature Medicine*, 6(11) 62-66, 2000) have indicated that autoimmune Encephalomyelitis, which is a model for MS, was ameliorated by AMPA/kainate which is an antagonist of the glutamate receptor.

Riluzole (2-amino-6-trifluoromethoxy benzothiazole) has recently emerged as a pharmacological agent potentially useful to slow down the evolution of

- 3 -

neurodegenerative diseases, such as amyotrophic lateral sclerosis. (Ben Simon *et al.*, *New Engl. J. Med.* 330:585-91 (1994)). In addition, this molecule has been shown to display anticonvulsant, anti-ischemic, and neuroprotective properties under various experimental conditions. A clear understanding of the site and
5 mechanism of action of this molecule is still lacking.

SUMMARY OF THE INVENTION

The present invention is based on the surprising findings that in an animal model of multiple sclerosis (MS) an experimental autoimmune encephalomyelitis (EAE) model produced by an injection of myelin oligodendrocyte glycoprotein
10 (MOG), animals treated by the administration of Riluzole before the appearance of clinical MS-related symptoms developed virtually no such clinical symptoms, and appeared essentially normal in all aspects tested. Moreover, treatment of such animals with Riluzole after the appearance of MS-related symptoms completely arrested the progression of the disease. Furthermore, some improvement in the
15 animal's condition had been shown.

The present invention provides, by one of its aspects, a pharmaceutical composition for the treatment of multiple sclerosis, comprising a pharmaceutically acceptable carrier, and, as an active ingredient, Riluzole.

The term "*treatment*" in the context of the invention refers to any one of the
20 following: amelioration of some of the undesired symptoms of multiple sclerosis; the prevention of the manifestation of such symptoms before they occur; slowing down or completely preventing the progression of the disease (as may be evident by longer periods between reoccurrence episodes, slowing down or prevention of the deterioration of symptoms etc.); enhancing the onset of the remission period;
25 slowing down the irreversible damage caused in the progressive-chronic stage of the disease (both in the primary and secondary stages); delaying the onset of said progressive stage, or a combination of two or more of the above.

Thus, it is clear that the pharmaceutical compositions of the invention are suitable to be administered to patients in all stages of the disease of multiple

- 4 -

sclerosis, including the initial relapsing-remitting stages, both during remission periods (to prevent or delay reoccurrence) or reoccurrence conditions (to expedite remission and delay the onset of the progressive stage), as well as in chronic-progressive stages.

5 The term "*multiple sclerosis*" refers also to any other autoimmune disease manifested by demyelination of the central nervous system's neurons. The first symptoms which appear at the onset of MS will be referred to herein at times as "*MS-related symptoms*". The symptoms of MS in EAE -induced animals (animal model of MS) are typically weakness and malfunction in the animal's tail, followed
10 by weakness of its rear feet and finally weakness in its front feet. In humans, such first MS-related symptoms may typically be double vision, facial numbness, facial weakness, vertigo, nausea, vomiting ataxia, weakness of the arms, etc.

 The term "*Riluzole*" refers to (2-amino-6-trifluoromethoxy-benzothiazole) and chemical modifications (such as addition, replacement or deletion of a
15 substituent) of the above. Those modifications which fall under the scope of the invention are those that feature the therapeutically effect of Riluzole (for example in the model specified in the "detailed description" part) at concentrations which are the same or lower than those of Riluzole.

 Typically, the pharmaceutical composition of the invention may be
20 administered to patients in any manner known in the art, such as parenteral administration, (intravenously, or intraparenteral), by injection or by oral administration, oral administration being preferred.

 The pharmaceutically acceptable carrier should be formulated in accordance with the intended mode of administration of the pharmaceutical composition, as
25 evident to any person versed in the art. For oral administration, the Riluzole may for example be formulated as a coated tablet comprising Riluzole as an active ingredient together with a pharmaceutically acceptable excipient such as, for example, calcium phosphate, microcrystalline cellulose, silica, magnesium stearate, polyethylene glycol, etc.

- 5 -

The dosage of the Riluzole in the pharmaceutical composition should be determined empirically for each individual, and should depend on various factors, such as the patients' age and weight, the stage of the multiple sclerosis (remitting-relapsing; chronic-progressive), the severity of the symptoms of each stage etc. For example only, the dosage of Riluzole which may be typically administered to an MS patient taken orally with the drug may be in the range of between about 50 mg to about 200 mg per day, preferably about 100 mg per day administered in two doses of 50 mg every 12 hours.

If desired, the pharmaceutical composition of the invention may further comprise other therapeutically effective agents which are known to be beneficial for the treatment of multiple sclerosis.

The present invention further concerns use of Riluzole for the preparation of a medicament for the treatment of multiple sclerosis.

By another aspect, the present invention concerns a method for treatment of multiple sclerosis, comprising: administering to a subject in need of such treatment, a therapeutically effective amount of Riluzole. Preferably the administration should be by oral administration.

BRIEF DESCRIPTION OF THE DRAWINGS

In order to understand the invention and to see how it may be carried out in practice, a preferred embodiment will now be described, by way of non-limiting example only, with reference to the accompanying drawings, in which:

Fig. 1 shows the mean clinical score of an EAE model of MS (mice injected with MOG) as a function of time after injection, for untreated mice (empty squares) and mice treated with Riluzole (full circles).

Fig. 2 shows the EAE clinical score model of MS (mice injected with MOG) as a function of time after injection, in mice treated with Riluzole (circles) and in untreated mice (squares).

- 6 -

Fig. 3 shows the EAE severity score model of MS (mice injected with MOG) as a function of time after injection, in mice treated with Riluzole at various times (circles, squares and triangles) and in untreated mice (rhombus squares).

DETAILED DESCRIPTION OF A PREFERRED EMBODIMENT

5 Example 1: Effect of Riluzole in an MS model.

Experimental Autoimmune Encephalomyelitis (EAE) induction:

EAE was induced in C3H.SW female mice (6 weeks old, Harlen, Rehovot, Israel) by immunization with the peptide encompassing amino acids 35-55 of rat myelin oligodendrocyte glycoprotein (MOG). Synthesis was carried out by the
10 Weizmann Institute Synthesis Unit, using a solid-phase technique on a peptide synthesizer (Applied Biosystems Inc., Foster City, CA City).

Mice were injected subcutaneously at one site in the flank with a 200 μ l emulsion containing 75 μ g MOG peptide in complete Freund's adjuvant (CFA) and 200 μ g Mycobacterium tuberculosis (Sigma Israel).

15

Treatment:

Mice (n=10) were injected i.p. with Riluzole 10mg/kg twice a day for 7 days before MOG injection and for additional 53 days after injection. Control mice (n=10) were injected twice a day with 200 μ l of saline for the same period.

20

Clinical score assessment:

0 = no clinical symptoms;

1 = loss of tail tonicity;

2 = partial hind limb paralysis;

25 3 = complete hind limb paralysis;

4 = paralysis of four limbs;

5 = total paralysis;

- 7 -

6 = death.

Results:

The results are shown in Fig. 1. Following the encephalitogenic challenge, mice were observed daily and clinical manifestations of EAE were scored. Four weeks after the encephalitogenic challenge, the saline injected mice (5/10) developed severe EAE characterized by complete hind limb paralysis (mean total score of 1.7 ± 0.2 SE) starting on day 36. In contrast, the Riluzole-treated mice were significantly resistant to MOG-induced EAE. Nine out of the ten immunized Riluzole-treated mice remained disease free with only one mouse demonstrated clinical signs only on day 54 ($p=0.00002$ using X2 values) with mean of total score of 0.1 ± 0.1 SE, ($p=0.015$ using Student-t-test).

Thus, Riluzole treatment markedly reduced both the incidence and clinical severity of the disease.

Example 2: Effect of Riluzole administered more than a week after the appearance of MS symptoms

EAE was induced in C3H.SW mice by injection of MOG as described in Example 1 above. The effect of administration of Riluzole on the progression of MS after the appearance of the first MS-related symptoms in the mice was tested.

Treatment:

Mice received MOG as described in Example 1 above and upon appearance of the first MS-related symptoms were divided into the following groups ($n=10$ in each group):

1. Mice receiving Riluzole 14 days after the appearance of the symptoms;
2. Mice which did not receive Riluzole (Control).

Riluzole was administered to the mice at a dose of 10mg/kg twice a day until the end of the experiment

The EAE score was calculated as described in Example 1 above.

Results:

As seen in Fig. 2, the first MS related symptoms were detected seven days after the injection of MOG. Mice were treated with Riluzole at day 14 and as seen
5 in the Fig., at day 15 there is a cessation in the development of MS in the treated mice with no alteration of the clinical score in these mice from this time on.

Example 3: Effect of Riluzole administered at various times, more than a week after the appearance of MS symptoms

10 EAE was induced in C3H.SW mice by injection of MOG as described in Example 1 above. The effect of administration of Riluzole on the progression of MS after the appearance of the first MS-related symptoms in the mice was tested.

Treatment:

15 Mice received MOG as described in Example 1 above and upon appearance of the first MS-related symptoms were divided into the following groups (n=10 in each group):

1. Mice receiving Riluzole upon the appearance of the symptoms, from day 12 to day 18;
- 20 2. Mice receiving Riluzole after the severe appearance of the symptoms (score=3.2 in an EAE model), from day 18 to day 28;
3. Mice which received Riluzole after the appearance of the symptoms on day 12, but stopped receiving Riluzole on day 18; and
4. Mice which did not receive Riluzole (Control).

25

Riluzole was administered to the mice at a dose of 10mg/kg twice a day until the end of the experiment (day 28).

The EAE score was calculated as described in example 1 above.

30

- 9 -

Results:

As seen in Fig. 3, sixteen days after the appearance of the first MS-related symptoms, mice that had been treated with Riluzole showed the lowest scores of EAE as compared to the untreated mice (0.9 ± 0.26 vs. 3.25 ± 0.48 , $p < 0.0004$).

5 Moreover, mice that started the Riluzole treatment in the late stage of the disease (on day 18) demonstrate significant improvement in their clinical score. Furthermore, Riluzole-treated mice that stopped the treatment on day 18 showed increase in the clinical score.

Thus, the results of these two experiments show that, in addition to the
10 effect of Riluzole administered to EAE-induced animals before they had developed first symptoms of the disease (Example 1), Riluzole treatment of EAE-induced animals shortly after the animals had developed first symptoms of the disease also markedly slowed down the progression of EAE and even improved the clinical manifestation.

- 10 -

CLAIMS:

1. A pharmaceutical composition for the treatment of multiple sclerosis (MS) comprising a pharmaceutically acceptable carrier and as an active ingredient, Riluzole.
- 5 2. A pharmaceutical composition for the treatment of autoimmune-demyelination of nerves comprising a pharmaceutically acceptable carrier and, as an active ingredient, Riluzole.
3. A pharmaceutical composition according to claims 1 or 2 for oral administration.
- 10 4. Use of Riluzole for the preparation of a medicament for the treatment of multiple sclerosis.
5. A method for the treatment of multiple sclerosis comprising administering to a subject in need of such treatment, a therapeutically effective amount of Riluzole.
- 15 6. A method according to claim 5, wherein the Riluzole is administered orally.
7. A pharmaceutical composition according to claims 1 or 2, wherein said composition is administered after the appearance of MS.-related symptoms.
8. A pharmaceutical composition according to claims 1 or 2, wherein the
20 treatment is administered before the appearance of MS-related symptoms.

1/3




FIG. 1

2/3





FIG. 2

3/3



Days after MOG injection

FIG. 3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IL 01/00534

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K31/428 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, MEDLINE, BIOSIS, SCISEARCH, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 00 74676 A (POLMAN CHRIS ; VERENIGING VOOR CHRISTELIJK WE (NL); BIOGEN INC (US)) 14 December 2000 (2000-12-14) the whole document	1-8
P, X	FERNANDEZ O.: "The rational basis of the newer treatments used in multiple sclerosis !. BASE RACIONAL PARA LOS NUEVOS TRATAMIENTOS EN LA ESCLEROSIS MULTIPLE." REVISTA DE NEUROLOGIA, (16 JUN 2000) 30/12 (1257-1264). , XP001041153 abstract; table 1	1-8

-/--

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z document member of the same patent family

Date of the actual completion of the international search

19 November 2001

Date of mailing of the international search report

26/11/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hoff, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IL 01/00534

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KALKERS N F: "A pilot study of riluzole in primary progressive multiple sclerosis; effect on spinal cord atrophy on MRI" AVAILABLE FROM INTERNET, 6 June 1999 (1999-06-06), XP002122164 the whole document	1-8
X	MCCREADY (ED) G: "Rilutek might be tried for MS" AVAILABLE FROM INTERNET, June 1998 (1998-06), XP002122163 the whole document	1-8
X	EP 0 558 861 A (RHONE POULENC RORER SA) 8 September 1993 (1993-09-08) the whole document	1-3,7,8
X	SCHLUEP M. ET AL: "Multiple sclerosis - Amyotrophic lateral sclerosis: Recent therapeutic progress!. NEUROLOGIE. SCLEROSE EN PLAQUES - SCLEROSE LATERALE AMYOTROPHIQUE: RECENT DEVELOPPEMENTS THERAPEUTIQUES." MEDECINE ET HYGIENE, (1997) 55/2145 (33-35). XP001041141 the whole document	1-3,7,8
A	STANKOFF B: "Neuroprotection and Central Nervous System myelination: New prospects for multiple sclerosis?" NEUROLOGY, LIPPINCOTT WILLIAMS & WILKINS, PHILADELPHIA, US, vol. 52, no. 6, SUPPL 2, 12 April 1999 (1999-04-12), page A402 XP002122165 ISSN: 0028-3878 the whole document	1-8

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IL 01/00534

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0074676 A	14-12-2000	AU 5421900 A WO 0074676 A1	28-12-2000 14-12-2000
EP 0558861 A	08-09-1993	FR 2688138 A1 AT 149833 T AU 666150 B2 AU 2948292 A CA 2117466 A1 CZ 9402120 A3 DE 69218255 D1 DE 69218255 T2 DK 627919 T3 EP 0558861 A1 EP 0627919 A1 ES 2098558 T3 WO 9317683 A1 GR 3022797 T3 HU 70946 A2 IL 103493 A JP 2713384 B2 JP 7504655 T KR 190296 B1 MX 9206109 A1 NO 943256 A RU 2110260 C1 SK 104794 A3 US 5527814 A ZA 9208213 A	10-09-1993 15-03-1997 01-02-1996 05-10-1993 16-09-1993 15-12-1994 17-04-1997 11-09-1997 12-05-1997 08-09-1993 14-12-1994 01-05-1997 16-09-1993 30-06-1997 28-11-1995 23-07-1996 16-02-1998 25-05-1995 01-06-1999 01-09-1993 02-09-1994 10-05-1998 10-05-1995 18-06-1996 30-04-1993

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.